


Ponderosa Pine





- Growth Form: conical to ovoid
- Crown Density: moderately dense
- Size: 40-100 feet high 15-60 foot spread
- Drought Resistance: excellent
- Cold Hardiness: excellent
- Growth Rate: rapid to moderate
- Life Span: long
- Elevational Range: to 9,000 feet
- Soil Conditions: tolerates alkaline well
- Possible Insect Problems: sawflies, bark beetle, tip moth
- Possible Disease Problems: western gall rust; dwarf mistletoe, shoestring root rot
- Wildlife Value: high: song and ground birds, small mammals, browsers
- Seasonal Color: evergreen
- Miscellany: native; needs well-drained soils



Ponderosa Pine

Pinus ponderosa



ZONE:	3-7	SHAPE:	Irregular
HEIGHT:	60-100'	EXPOSURE:	Full Sun
SPREAD:	25-30'	GROWTH RATE:	Medium
SOILS:	Will grow on wide range of soils – alkaline, loamy, moist, well-drained, and very sandy soils with little topsoil. Once established, is very drought-tolerant.		

DESCRIPTION:

- Rugged-looking. Lean. Enduring. Stately. If a tree could personify the West, it would surely have to be the ponderosa pine. It is also one of America's most abundant tree species. It is a mainstay in the lumber industry but has also been put to work in windbreaks, buffer strips, reclamation and mass landscape plantings. Its size, fast growth, green foliage, interesting cinnamon-colored bark texture, pleasant aroma and large ornamental cones also make for a popular specimen or ornamental tree. The mature thick bark helps to make it fire resistant, and its deep taproot very wind resistant. It is long lived - up to hundreds of years old.

WILDLIFE VALUE:

- The seeds provide food for birds and small mammals, particularly turkeys, nuthatches, crossbills, grosbeaks, pine siskins, grouse, squirrels, chipmunks, and mice. The leaves, twigs, and bark are browsed by porcupines, mule deer, and elk. Snags provide a large number of wildlife species with nesting and roosting sites.

HISTORY / LORE / USE:

- It is one of the most widely distributed pines in western North America. It occupies diverse habitats throughout the west from southern British Columbia to Mexico, and Nebraska and Oklahoma to the Pacific coast. The Scottish botanist David Douglas named this pine for its ponderous or heavy wood. Ponderosa pine forests are valuable for timber production, livestock grazing, recreation, and wildlife habitat. Native Americans used the inner bark for emergency flour and boiled the young cones for emergency food. In the spring the bark was scraped and eaten raw as a sweet treat. Inner bark gum was used for medicine. The needles were steeped to make a tea.

MOISTURE:

- The ponderosa pine prefers moist, well-drained soil, but will grow in dry, arid areas and is highly drought tolerant when well established. It is intolerant of excessively wet sites or sites with poor drainage.

LEAVES:

- The needles are sharp, stiff but flexible, 5-10" long, densely crowded on the branchlets in bundles of three (sometimes two), dark-gray green or yellowish green. They persist on the tree for 3-5 years and then turn brown and are shed.

FLOWER COLOR:

- Male flower is yellow, female is red purple in pairs.

BLOOM TIME:

- Spring

FRUIT DESCRIPTION:

- Cones are solitary or in groups of 3 to 5. They are 3-6" long, light reddish brown, matte or glossy. May be produced as early as seven years with irregular good crops depending upon the variety.

PONDEROSA PINE

Pinus ponderosa P. & C.

Lawson

Plant Symbol = PIPO

Contributed by: USDA NRCS National Plants Data Center



Banner, R. 2002.
Utah State University Extension.

Alternate Names

Big heavy, black jack, bull pine, ponderosa white pine, Sierra brown bark pine, silver pine, western pitch pine, western red pine, western yellow pine, yellow pine, Yosemite pine.

Uses

Erosion control: Ponderosa pine is a rapid growing tree with the ability to firmly anchor into most soil types. For this reason, it is suitable for use as a windbreak species. It can also be used with other natives to provide cover and erosion control on rehabilitated sites.

Ethnobotanic: Native Americans used various parts of ponderosa pine for medicinal, building and household, food, and ceremonial purposes. Needles were used as dermatological and gynecological aids.

They were also used to reduce coughs and fevers. The pitch was used as an ointment for sores and scabby skin, backaches, rheumatism, earaches, inflamed eyes, and as a sleeping agent for infants.

The boughs of the plant were used in sweat lodges for muscular pain, as decoctions for internal hemorrhaging, and as infusions for pediatric treatments.

The roots of ponderosa pine were used to make blue dye and needles were used as insulation for underground storage pits. The wood was used extensively for fence posts, boards for general construction, and to fabricate snowshoes. Single logs were used to make dugout canoes. Bark was used to cover houses.

Most parts of the plant were used for food, including the pitch, seeds, cones, bark, buds, and cambium. The pollen and needles were used in healing ceremonies.

Ornamental value: Ponderosa pine has a lush green color and pleasant odor that makes it popular for ornamental plantings. It has been planted, sometimes out of its natural range, because of its aesthetic qualities. Ponderosa pine is used as borders of forested highways, but is not planted within the right-of-way. The large stature of the tree limits its use to open spaces.

Wildlife: Red-winged blackbirds, chickadees, mourning doves, finches, evening grosbeak, jays, Clark's nutcracker, nuthatches, rufous-sided towhee, turkeys, chipmunks and squirrels consume the seeds of ponderosa pine. Blue and spruce grouse use ponderosa pine needles for nesting material. Mice, porcupines, and other rodents use the bark for nesting material. The trees are also important to various birds for cover, roosting and nesting sites.

Wood production: Ponderosa pine is one of the most important timber species in the western United States. The annual production of ponderosa pine is ranked third behind Douglas fir and hem-fir. Approximately 1.3 billion board feet of ponderosa pine lumber is produced annually out of Oregon, the largest supplier in the United States. It is popularly used for the construction of buildings.

Description

General: Pine Family (Pinaceae). Ponderosa pine is a large tree that lives 300 to 600 years and reaches heights of 30 to 50 m tall and 0.6 to 1.3 m in diameter. The oldest trees can exceed 70 m in height and 2 m in diameter. The bottom one-half of the

straight trunk is typically without branches. The crown of ponderosa pine is broadly conical to round-shaped. The bark is characteristically orange-brown with a scaly plate-like appearance. Twigs are stout, up to 2 cm thick, orange-brown, and rough. Needles are 12 to 28 cm long, thin and pointed with toothed edges, occur in bundles of three, and give a tufted appearance to the twig. Buds are up to 2 cm long, 1 cm wide, red-brown with white-fringed scale margins. Male cones are orange or yellow and are located in small clusters near the tips of the branches. The female cone is oval, woody, 8 to 15 cm long, with a small prickle at the tip of each scale. Flowering occurs from April to June of the first year, and cones mature and shed winged seeds in August and September of the second year.

Distribution: Ponderosa pine is distributed from southern British Columbia through Washington, Oregon, and California, and east to the western portions of Texas, Oklahoma, Nebraska, North Dakota, and South Dakota. For current distribution, please consult the Plant Profile page for this species on the PLANTS Web site (<http://plants.usda.gov>).

Habitat: Ponderosa pine trees occur as pure stands or in mixed conifer forests in the mountains. It is an important component of the Interior Ponderosa Pine, Pacific Ponderosa Pine-Douglas fir, and Pacific Ponderosa Pine forest cover types.

In the northwest, it is typically associated with Rocky Mountain Douglas fir, lodgepole pine, grand fir, and western larch. In California it is associated with California white fir, incense cedar, Jeffrey pine, sugar pine, coast Douglas fir, California black oak, and western juniper. In the Rocky Mountains and Utah, it is associated with Rocky Mountain Douglas fir, blue spruce, lodgepole pine, limber pine, and quaking aspen. In the Black Hills, it is associated with quaking aspen, white spruce, and paper birch. In Arizona and New Mexico, it is associated with white fir, Rocky Mountain Douglas fir, blue spruce, quaking aspen, gamble oak, and southwestern white pine at higher elevations and Rocky Mountain juniper, alligator juniper, and Utah juniper at lower elevations (Oliver & Riker 1990).

Shrubs and grasses typically associated with ponderosa pine within its range include ceanothus, sagebrush, oak, snowberry, bluestem, fescue, and polargrass.

Adaptation

The USDA hardiness zones for ponderosa pine range from 3 to 7. It grows on a variety of soils from

shallow to deep, and from gravelly sands to sandy clay loam. It is found growing on bare rock with its roots in the cracks and crevices. It has a low tolerance to alkalinity, preferring soils with a pH of 6.0 to 7.0. It grows best in zones with 30 to 60 cm average annual precipitation on well-drained soils. Once established it also survives hot and dry conditions, exhibiting medium to good drought tolerance. Fifty percent shade reduces the growth rate significantly. It withstands very cold winters.

Ponderosa pine is a climax species at the lower elevations of the coniferous forest and a mid-successional species at higher elevations where more competitive conifers are capable of growing. It generally grows at elevations between sea level and 3,000 m. The populations at higher elevations usually occur within the southern part of its range (Oliver & Riker 1990).

Establishment

Site preparation is needed to control competition, which compromises seedling survival and growth. Seeds are sown in late March to early April. The seed is sown for an initial density of 237 seedlings/m² (22 seedlings/ft²). Transplant stock should be one or two years old, with less than 2 prior transplantings, and 15 to 30 cm in height. Space the plants 1 to 3 m apart depending on the site.

Initial seedling survival is reduced under moisture stress. Older seedlings can tolerate limited moisture. Competition from other vegetation should be controlled for the first three to six years until the trees become well established.

Management

Ponderosa pine can be over-irrigated in poorly drained soils, or drowned out on high water table sites.

It responds well to thinning, which should be done as stands become older to develop larger crowns, resulting in heavier seed crops for wildlife. More forage for deer and elk become available from associated plants by opening the canopy. The use of repellents or other control measures may be necessary to prevent overuse of the trees by rodents.

Ponderosa pine is resistant to fire due to its thick bark. Low intensity surface fires control competitive species like scrub oak and shade-tolerant conifers. Ponderosa pine seedlings can also survive low intensity burns.

Pests and Potential Problems

Approximately 200 insect species affect ponderosa pine from its cone stage to maturity. Pine cone beetles cause tree death by transmitting blue stain fungus to the tree. Their larvae also consume the phloem, restricting the flow of nutrients to the top of the tree.

Western pine beetle is a common cause of death for older trees, drought stressed trees, and even healthy, vigorous trees during epidemics.

Bark beetles are naturally present in all stands. Harvesting methods that leave large amounts of logging slash can allow bark beetle populations to explode and kill vigorous trees up to 0.5 m in diameter.

The ponderosa pine budworm, also known as the sugar-pine tortrix, eats new needles on trees in New Mexico and Colorado. Several years' worth of damage will affect the health of the tree. Early research suggests that some insecticides may help to control infestations.

Dwarf mistletoe is the most widespread parasite that causes branch and stem deformation. It germinates on ponderosa pine branches and forces its roots into the phloem of the host branch, creating stem cankers that leave the wood weak and unsuitable for use as lumber. This weakens the tree and leaves it susceptible to fungal infections and insect attacks. Root diseases, rusts, trunk decays, and needle and twig blights also cause significant damage.

Seeds and Plant Production

Ponderosa pine is propagated by seed. Cones are ready for collection in October and November when they turn reddish brown. Mature seed is firm and brown in color. Cones should be dried on canvas tarp in a well-ventilated area immediately after they have been collected. The seeds will drop from the cones as they dry.

Several germination methods for ponderosa pine have been utilized, each with their own variations. In general, seeds undergo an imbibition treatment before stratification. Seeds are placed in mesh bags and soaked in cold running water for 48 hours. One variation is to soak the seeds in a 40% bleach solution for 10 minutes with hand agitation prior to placing them under running water. The mesh bags are placed in plastic bags and stored at 1°C for 2 to 8 weeks. They should be checked daily for mold. Seeds are sown into containers and covered with media. The media should be kept moist throughout

germination. Germination will occur at an average greenhouse temperature of 20°C. Alternating greenhouse temperatures of 21-25°C during the day and 16-18°C at night is an appropriate environment for germinating seeds. Germination will occur in approximately 15 days.

Seedlings are thinned and watered daily throughout the establishment phase. They should not be moved outdoors until after the last frost of the year.

Seeds can be dried to between 5 and 8% moisture and placed in airtight plastic bags, then stored for long periods of time in freezers set at -15°C.

Cultivars, Improved and Selected Materials (and area of origin)

Contact your local Natural Resources Conservation Service (formerly Soil Conservation Service) office for more information. Look in the phone book under "United States Government." The Natural Resources Conservation Service will be listed under the subheading "Department of Agriculture."

References

- Agricultural Research Center. 2004. *GRIN taxonomy* (<http://www.ars-grin.gov/cgi-bin/npgs/html/index>, 27 May 2004). USDA, Beltsville.
- Banner, R. 2002. *Ponderosa pine* (<http://extension.usu.edu/rangeplants/Woody/ponderosa.htm>, 29 July, 2004). Utah State University Extension, Logan.
- Forest Research Nursery. 2004. *Native plant network* (<http://www.nativeplantnetwork.org>, 27 May 2004). College of Natural Resources, University of Idaho, Moscow.
- Frankis, M.P. 1998. *Pinus ponderosa*. Earle, C.J., *Gymnosperm database* (<http://www.conifers.org/>, 27 May 2004). Bonn University, Bonn, Germany.
- Miller, A. 1997. *Ponderosa pine species facts* (<http://www.wvpa.org/ppine.htm>, 16 June 2004). Western Wood Products Association, Portland.
- Moerman, D. 2003. *Native American ethnobotany database* (<http://herb.umd.umich.edu/>, 27 May 2004). University of Michigan, Dearborn.
- Oliver, W.W. and R.A. Ryker. 1990. *Ponderosa pine*. Burns, R.M. and B.H. Honkala, *Silvics of North America. Agriculture Handbook 654*. USDA Forest Service, Washington, D.C.

Rewerts, M.S. 2004. *Ponderosa pine budworm*
(<http://www.ext.colostate.edu/pubs/insect/05567.html>
, 29 July, 2004). Colorado State University
Extension, Fort Collins.

Prepared By:

Sarah Wennerberg
Formerly USDA NRCS National Plant Data Center,
Baton Rouge, Louisiana

Species Coordinator:

Mark Skinner
USDA NRCS National Plant Data Center, Baton
Rouge, Louisiana

Edited: 29Jul2004 sbw; 20Oct2004 rln; 060802 jsp; 060830 jsp

For more information about this and other plants, please contact
your local NRCS field office or Conservation District, and visit the
PLANTS Web site<<http://plants.usda.gov>> or the Plant Materials
Program Web site <<http://Plant-Materials.nrcs.usda.gov>>

*The U.S. Department of Agriculture (USDA) prohibits
discrimination in all its programs and activities on the basis of
race, color, national origin, sex, religion, age, disability, political
beliefs, sexual orientation, and marital or family status. (Not all
prohibited bases apply to all programs.) Persons with disabilities
who require alternative means for communication of program
information (Braille, large print, audiotape, etc.) should contact
USDA's TARGET Center at 202-720-2600 (voice and TDD).*

*To file a complaint of discrimination write USDA, Director, Office
of Civil Rights, Room 326-W, Whitten Building, 14th and
Independence Avenue, SW, Washington, DC 20250-9410 or call
202-720-5964 (voice or TDD). USDA is an equal opportunity
provider and employer.*

*Read about Civil Rights at the Natural Resources Conservation
Service.*